

Published in final edited form as:

Clin Cancer Res. 2010 August 1; 16(15): 3976–3987. doi:10.1158/1078-0432.CCR-09-3358.

Genetically Abnormal Circulating Cells in Lung Cancer Patients: An Antigen Independent Fluorescence in-situ Hybridization Based Case-Control Study

Ruth L. Katz¹, Weigong He¹, Abha Khanna¹, Ricardo L. Fernandez¹, Tanweer M. Zaidi¹, Matthew Krebs⁷, Nancy P. Caraway¹, Hua-Zhong Zhang¹, Feng Jiang⁸, Margaret R. Spitz², David P. Blowers⁶, Carlos A. Jimenez⁴, Reza J. Mehran³, Stephen G. Swisher³, Jack Roth³, Jeffrey S. Morris⁵, Carol Etzel², and Randa El-Zein²

¹ Department of Pathology, the University of Texas M. D. Anderson Cancer Center, Houston, TX

² Department of Epidemiology, the University of Texas M. D. Anderson Cancer Center, Houston, TX

³ Department of Thoracic and Cardiovascular Surgery, the University of Texas M. D. Anderson Cancer Center, Houston, TX

⁴ Department of Pulmonary Medicine, the University of Texas M. D. Anderson Cancer Center, Houston, TX

⁵ Department of Biostatistics, the University of Texas M. D. Anderson Cancer Center, Houston, TX

⁶ AstraZeneca, Macclesfield, Cheshire, UK

⁷ Paterson Institute for Cancer Research, Manchester, UK

⁸ University of Maryland School of Medicine, Baltimore, MD

Abstract

Purpose—We performed a study to determine if a *fluorescence in-situ hybridization* (FISH)-based assay using isolated peripheral blood mononuclear cells (PBMCs) with DNA probes targeting specific sites on chromosomes known to have abnormalities in Non Small Cell Lung Cancer (NSCLC) cases could detect circulating genetically abnormal cells (CACs).

Experimental Design—We evaluated 59 NSCLC cases with stage I through IV disease and 24 controls. PBMCs and matched tumors were hybridized with 2 two-color (3p22.1/CEP3 and 10q22.3 [SP-A]/CEP10) and 2 four-color (CEP3, CEP7, CEP17, and 9p21.3 [URO]) and (EGFR, c-MYC, 6p11-q11, and 5p15.2 [LAV]) FISH probes. Percentages of cytogenetically abnormal cells (CACs) in peripheral blood and in matched tumor specimens were quantified using an automated fluorescent scanner. Numbers of CACs were calculated based on the percentage of CACs (defined as PBMCs with genetic abnormalities) per mL of blood and expressed per microliter of blood.

Results—Patients with NSCLC had significantly higher numbers of CACs than did controls. Mean number of CACs ranged from 7.23±1.32/μl for deletions of 10q22.3/CEP10 to 45.52±7.49/μl for deletions of 3p22.1/CEP3. Numbers of CACs with deletions of 3p22.1, 10q22.3, and 9p21.3, and gains of URO, increased significantly from early to advanced stage of disease.

Conclusions—We have developed a sensitive and quantitative antigen-independent FISH-based test for detecting CACs in peripheral blood of patients with NSCLC which showed a significant correlation with the presence of cancer. If this pilot study can be validated in a larger study, CACs may have a role in the management of patients with NSCLC.

INTRODUCTION

Cytogenetic evidence exists to show that circulating epithelial cells (CECs) in the bloodstream contain genetic abnormalities similar to those in primary tumors (1). Investigators employed a combination of immunohistochemistry and *fluorescence in-situ hybridization* (FISH) to demonstrate that the aneusomies found in CECs were similar to those detected in primary breast, colon, kidney and prostate cancers (1). Detection of CECs or circulating tumor cells (CTCs) using a simple blood test may assist in early detection of lung cancer at diagnosis and relapse and provide a minimally invasive way to monitor results of therapy. Researchers have attempted to isolate CTCs from peripheral blood using a variety of techniques, including immunomagnetic capture of EpCAM-positive cells, which have demonstrated to be of clear prognostic use for breast, prostate, and colorectal carcinomas (1–7). However, usually very few CTCs are recovered using these techniques. Other CTC isolation techniques include filtration of CTCs, density gradient separation, microchip technologies and polymerase chain reaction (PCR) based techniques (8–12). However, there are limitations to these methods, particularly PCR which restricts enumeration and morphological analysis of cells (13–14).

Relatively few reports have described detection of CTCs in patients with lung cancer. A recent report described positive depletion of CD45+ cells from blood with magnetic beads followed by CK18/19 staining of enriched cells in patients with non-small cell lung cancer (NSCLC) (15). While few CTCs from these patients were recovered, the number of CTCs generally correlated with the stage of disease, with fewer than two CTCs per 7.5 mL of blood detected in controls. However, relying on expression of epithelial markers in CTCs, including lack of sensitivity to antigenic detection by EpCam, may drastically reduce the sensitivity of CTC detection if the epithelial markers are absent. For example, a recent study pointed to the expression of mesenchymal rather than epithelial markers in common types of breast carcinoma, which may have resulted in undercounting of CTCs (9).

The use of a FISH assay without an epithelial capture component to detect genetically abnormal circulating cells is attractive, as it does not rely on antigenic expression and may reveal more CACs than anticipated, if not limited by an antigenic assay for epithelial differentiation. FISH-based tests detect losses and gains of selected chromosomal regions in the nuclei of cells regardless of the lineage of these cells, and can be used to simultaneously evaluate up to four different DNA probes in a single cell. Rapid automated fluorescent scanning of cells and/or analysis of multicolor fluorescent signals for different genes on a per-cell basis using FISH have enabled the classification and quantitation of cytogenetically abnormal cell populations (CACs) in urine, blood and sputum (16–19). Based on a previous study of lung adenocarcinomas and squamous cell carcinomas using CGH arrays we selected two genes that are frequently deleted in NSCLC, 3p22.1 (GC20) and 10q22.3 (surfactant protein A1 and A2, SP-A) and verified that they were deleted in NSCLC specimens by FISH (18). We previously demonstrated more frequent deletions of these genes in sputum specimens obtained from patients with NSCLC than in controls using FISH (18).

Based on these previous results, in the present study we sought to establish whether CACs can be detected in the bloodstream of patients with NSCLC. We hypothesized that the CACs encountered in the circulation would bear similar genetic abnormalities to those detected in NSCLC. We used four sets of genome-specific FISH probes, including 3p22.1/CEP3, 10q22.3/CEP10, and two commercially available probe sets that were selected based on genetic profiles

of genes known to be important in the pathogenesis of lung and bladder cancer (20–36) to test peripheral blood mononuclear cells (PBMCs) obtained from healthy controls without lung cancer (smokers and nonsmokers) and from patients with NSCLC at different stages. We also tested resected tissue specimens from corresponding primary lung tumors using the same panel of biomarkers.

The main objective of this study was to elucidate and compare the number of CACs between patients with NSCLC and controls, to determine whether these cells, based on their genetic abnormalities, were indicative of metastasis from the primary tumor. Secondary aims included correlating the presence of number of CACs in the blood with relapse rate and survival duration; comparing the genetic abnormalities in CACs in the blood with those in the corresponding lung tumors, and developing a quantitative assay for CACs per microliter of blood based on the percentages of CACs in peripheral blood.

Material and Methods

Study Population

In 2007 and 2008, peripheral blood specimens were collected prospectively from 59 patients with NSCLC and 24 controls including heavy smokers at high risk for lung cancer at The University of Texas M. D. Anderson Cancer Center under an Institutional Review Board-approved protocol. High risk controls were patients enrolled in a lung cancer screening trial. Cases and controls were not matched by age, smoking status or sex as seen in Table 1. However, we adjusted for these variables in the analysis.

Demographic Characteristics of the Study Population

Criteria for study entry included no treatment prior to surgery for stage I–III NSCLC cases. Equal stratification of patients across all NSCLC stages was attempted. Corresponding primary lung tumor tissue specimens were available for 21 patients. Mean ages of the controls and patients were 55.5 ± 2.86 and 66.8 ± 1.36 years, respectively (Table 1). Disease stages ranged from early (14 IA, 8 IB, and 9 II) to advanced (10 III and 18 IV), and adenocarcinoma was the most common subtype. Of 23 patients who had a relapse or persistent disease, 4 had an early relapse (within 6 months to 1 year after first treatment). At the time of data analysis, 22 patients had died, most of whom had stage III or IV disease.

Specimen Collection

Twenty-five mL of blood in ethylenediaminetetraacetic acid (EDTA) was collected from controls, patients with operable disease immediately before surgery, and patients with inoperable disease, at the time of endoscopic ultra-sound bronchoscopic evaluation performed for staging. Following collection, blood was immediately processed by Ficoll-Hypaque density gradient separation. Peripheral blood mononuclear cells (PBMCs) were isolated and counted using a Coulter counter (Beckman Coulter, Fullerton, CA), and cytospin preparations of PBMCs containing an average of 10,000 cells were prepared, spray-fixed with alcohol and stored at -20°C until needed. PBMCs from patients and controls were tested with the same panel of biomarkers. Tumor tissue was available on 21 patients who were enrolled in a lung cancer Specialized Program of Research Excellence study. FISH was performed on both the peripheral blood and tumor tissue to detect concordance of genetic abnormalities in both surrogate and target tissues.

FISH Testing for Cytogenetically Abnormal PBMCs (CACs)

The following panel of FISH probes was used: 1) a combination of two probe sets: Locus Specific Identifier (LSI) 3p22.1 with corresponding centromeric probe CEP3 and LSI 10q22.3

[SP-A] with corresponding CEP10 prepared in-house as described previously (18,21,22,23) and 2) two commercially available probe sets containing four probes each - LAVysion [LAV]: *EGFR*, *C-MYC*, 6p11-q11, and 5p15.2; and UroVysion [URO]: CEP3, CEP7, CEP17, and p16^{INK4A} 9p21.3 (Abbott Molecular, IL). Fluorescent signals in specimens were quantitated on a per-cell basis using an automated fluorescent system (Bioview, Rehovoth, Israel) that is capable of scanning and classifying hundreds of cells under fluorescent illumination and allows for detection of rare cells according to FISH pattern (19). Using two-color FISH with 3p22.1/CEP3 and 10q22.3/CEP10 a mean of 250 PBMCs was accumulated for each probe set and reviewed for appropriate morphology (round or oval cells) and to verify the number of FISH signals displayed by the program on a per-cell basis by an experienced observer blinded to the disease status. Similarly, at least 200 PBMCs were selected and scored for genomic abnormalities using both URO and LAV four-color probe sets. Cytogenetic abnormalities were scored based on the presence of chromosomal deletions, gains, monosomy, polysomy, or the sum of all abnormalities combined and expressed as percentages of CACs (for detailed methods of scoring biomarker abnormalities see supplemental material).

CAC Quantitation

The number of CACs per microliter of blood was calculated as the percentage of CACs (for a specific chromosomal probe set) \times the total number of PBMCs isolated/mL of blood collected/1000. Thus, the number of CACs with deletions or gains of 3p22.1 compared with CEP3 and the number of CACs with deletions or gains of 10q22.3 compared with CEP10 per microliter were calculated. CACs per microliter were calculated for the URO and LAV probe sets based on the presence of at least two chromosomal abnormalities in the biomarkers tested in each nucleus.

Tumor Wash Specimens

Cell suspensions of tumors were available from 21 patients with resectable tumors and cytopins of tumor cells were prepared. FISH was performed for the 3p22.1/CEP3, 10q22.3/CEP10, the URO and LAV probe sets and evaluated as detailed in the supplementary materials. A subset of tumors was further classified according to clonal heterogeneity based on chromosomal abnormalities for the different DNA biomarkers (Table 6S).

Recovery of Lung Adenocarcinoma Cell Line Experiments

The sensitivity of the FISH-based assay to detect the presence of CACs in peripheral blood was evaluated by performing recovery experiments in which H1299 lung adenocarcinoma cells (AC) were spiked into PBMCs isolated from healthy donors. Four separate dilution assays at 1 AC/10,000, 1 AC/1000, 1 AC/100 and 5 AC/100 were performed. The spiked cell mixtures were hybridized with 3p22.1/CEP3, 10q22.3/CEP10, the LAV set, and the URO set. H1299 cells and PBMC controls were similarly hybridized and evaluated for cytogenetic abnormalities. Calibration curves depicting the number of cells recovered for each biomarker at the different serial dilutions are shown in Figure 1 and Table 1S.

Statistical Analysis

Descriptive statistical analyses, including the Spearman χ^2 test, were used to test for distributional differences between the patients and controls according to categorical variables, and the Mann-Whitney test was used to determine differences in continuous variables. The Mann-Whitney test was also used to test for differences in each biomarker between the patients and controls. Simple linear regression analysis was performed to test for trends in the biomarkers by disease stage. Two-sided *P* values were used to determine the level of significance for each test.

To evaluate the role of each biomarker in cancer recurrence and overall survival we dichotomized each variable into two groups based on the 75th percentile of the controls for each respective outcome. Time to recurrence was defined as the number of months from the date of first treatment to that of first recurrence. Overall survival time was defined as the number of months from the date of first treatment to that of death. Patients lost to follow-up or those patients who had no recurrences or did not die prior to the end of the study were censored. The Kaplan-Meier method was used to identify any significant differences in time to recurrence and overall survival between the high and low groups for each biomarker, respectively. The time to recurrence and overall survival between the high and low biomarker groups was compared using the log-rank test. Biomarkers found to be significant at the 10% level in the log-rank test were further evaluated using the Cox proportional hazards model adjusted for age, sex, and disease stage.

RESULTS

Cell Recovery Validation Study

The rate of identification of H1299 carcinoma cells based on cytogenetic abnormalities exceeded 99%, whereas unspiked PBMCs with more than 1% chromosomal abnormalities were not identified. Tumor cell-recovery rates for H1299 using polysomy of CEP3/3p22.1 and 10/10q22.3 DNA probes to quantitate tumor cells at 0.01%, 0.1%, 1.0%, and 5.0% dilution levels ranged from 0.011%, 0.102%, 0.4% to 3.6% respectively for the CEP3/3p22.1 probe set, and similar recovery rates noted for the CEP10/10q22.3 (Figure 1 and Table 1S). For URO and LAV probe sets tumor cells were counted only if two abnormal chromosomes per nucleus were present and actual recovery of H1299 carcinoma at 0.01% to 5.0% dilutions ranged from 0.008% to 5.0% for the LAV set and 0.01% to 5.0% for the URO probe set. (Figure 1 and Table 1S). These results demonstrated that the assay recovered the correct proportion of abnormal cells throughout the range of the dilutions from 1/20 (5%) down to 1/10000 (0.01%).

CACs by Case, Control Status and NSCLC Stage

We recorded the mean percentage of circulating CACs in cases and controls and stratified them by pathological stage or clinical stage in cases that were inoperable because of advanced disease (IIIB or IV). There was a significant trend for percentages of CACs for all biomarkers, except the LAV probe set, to increase from early stage to advanced stage disease. We noted highly significant differences in the biomarker distribution between the patients and controls (Table 2A). For example, the mean \pm standard error of the mean [SEM] percentage of CACs in the controls ranged from 0.17 ± 0.07 for 3p22.1/CEP3 gains to 3.05 ± 0.46 for combined 3p22.1/CEP3 chromosomal abnormalities. In comparison, the mean \pm SEM percentage of CACs for patients with stage IA NSCLC ranged from 1.11 ± 0.30 for 3p22.1/CEP3 gains to 7.00 ± 0.93 for combined 3p22.1/CEP3 abnormalities. Both EGFR deletions and gains were significantly different between cases and controls (Figure 2A and Table 2A).

The mean numbers \pm SE of CACs per microliter (derived from the percentages of CACs) are depicted according to the stage of disease and cases were significantly different compared to controls for all biomarker abnormalities recorded (Figure 2B and Table 2B). Expressed per milliliter the mean number of CACs for all cases of NSCLC ranged from $7,230 \pm 1320$ for deletions of 10q22.3/CEP10 to $45,520 \pm 7490$ for deletions of 3p22.1/CEP3, while for URO and LAV abnormalities mean CACs were $18,790 \pm 3160$ and 17570 ± 2820 respectively.

Biomarker Abnormalities Associated with Tumor Stage

Many CACs were significantly associated with early-stage IA and/or advanced stage (IIIA, IIIB, or IV) NSCLC ($P < 0.05$) [Supplemental Table 3S]. Most notable were CACs containing abnormalities of 3p22.1/CEP3 and 10q22.3/CEP10, and gain or loss of biomarkers in the URO

set which increased significantly from early to advanced stage disease (Supplemental Figure 1S).

Correlation between Blood and Corresponding Lung Cancer Tissue

We obtained paired sets of peripheral blood and tumor tissue from 21 patients who underwent surgical resection of their lung tumors. The same set of FISH probes was used in both the PBMCs and tumor specimens. We observed a strong overall correlation between eight biomarker abnormalities in PMBCs and corresponding biomarkers in the tumor washes; specifically, six were positively correlated and included gains of *EGFR*, *C-Myc*, 6p11-q11, 3p22.1 and different abnormalities in the URO set (Table 5S). Overall, *EGFR* gain in CACs was significantly correlated with *EGFR* gains in tumor washes, especially among patients who presented with stage III and IV disease ($P \leq .01$). We observed positively correlated chromosomal abnormalities in the CACs and those in the tumor cells by the URO probe set. In contrast, the genetic abnormalities in the LAV set in CACs were negatively correlated with those in the tumor washes. An example of CACs and corresponding tumor cells for all biomarkers quantitated is depicted in Figures 3A and 3B.

Biomarker Abnormalities Associated with Disease Recurrence

Twenty-three (39%) patients had disease recurrence. The median time to recurrence was 29 months [95% confidence interval (CI), 15.49 to 42.51 months]. Twenty biomarkers were significant at the 10% level, of which, twelve were significant at 5% level in Kaplan-Meier analyses for disease recurrence (Figure 2S, Table 3). Of these, three were significant at the univariate level using the Cox proportional hazards model: namely 5p15.2 gain, 3p22.1 deletion, and a single URO gain. However, these biomarkers were not significant for disease recurrence after adjustment for age, sex, and disease stage.

Biomarker Abnormalities Associated with Overall Duration of Survival

Twenty-two (37%) of the patients died over a period of less than 1 month to 39 months following collection of baseline blood samples. The median overall survival duration was 29 months (95% CI, 12.69 to 45.31 months). Six biomarkers were significant at the 10% level in Kaplan-Meier analyses for overall survival (Figure 3S) but only two were significant at the univariate level: *EGFR* deletions and a single URO gain. However, these biomarkers were not significant for overall survival after adjustment for age, sex, and disease stage (Table 3).

DISCUSSION

To our knowledge, this is the first study to use a quantitative FISH-based assay to detect cytogenetically abnormal circulating cells from patients with stage (I–IV) NSCLC and controls, using specific probes to identify previously characterized genomic aberrations described in NSCLC. We hypothesized that CACs encountered in the circulation would bear similar genetic abnormalities to those detected in NSCLC. However, in order to prove that these CACs are of epithelial origin, current standards require demonstration of expression of epithelial markers such as cytokeratin or EpCAM, with negative expression of CD45, a hematolymphoid marker. Alternatively if CACs derived from NSCLCs have undergone an epithelial to mesenchymal transition (EMT) antigenic expression of vimentin, SNAIL, TWIST1 or other markers of EMT with concurrent demonstration of genetic abnormalities by FISH would be required (37). If circulating cancer stem cells with cytogenetic abnormalities are present, demonstration of ALDH1 or CD133 in these cells would be needed (38).

In addition it is possible that other cells present in the circulation including endothelial precursor cells, lymphocytes, monocytes, and hematopoietic stem cells may express aberrant genetic markers. Cytogenetic abnormalities have previously been demonstrated in peripheral

blood lymphocytes of patients with lung cancer that were identical with those of the primary tumor (39).

Previous investigators used FISH to detect aneuploidy in CTCs obtained by immunomagnetic capture for epithelial antigen followed by immunohistochemistry for cytokeratin or prostate specific antigen (40,41). While these studies, demonstrated genetic abnormalities similar to those of the primary tumor, they were limited by a low cell recovery and inability to detect chromosomal abnormalities in patients with <10 CTCs per 7.5 mL blood (17,40,41). To detect CACs we used an automated fluorescence-based scanner to analyze a minimum of 900 PBMCs obtained by density gradient enrichment and hybridized for 12 different DNA biomarkers in all patients and controls. While we hesitate to call these CACs “circulating epithelial cells” based on lack of phenotypic evidence for epithelial cell differentiation, we assume that the CACs detected may in fact be CTCs derived from NSCLC based on the following findings: 1) eight of the DNA biomarkers in the PBMCs were significantly correlated with the resected lung tumors; 2) the mean percentages of genetically altered cells for all 12 biomarkers tested correlated with the stage of NSCLC, with the lowest levels detected in patients with stage I disease and the highest detected in patients with stage III and IV disease; 3) the controls had significantly fewer genetically abnormal cells for all the biomarkers; 4) at the 0.0001 dilution level of spiked tumor cells in PBMCs from buffy coats of normal healthy donors, genetic abnormalities, such as polysomy for CEP10/10q22.3 or CEP3/3p22.1 genes or abnormal cells by FISH for either of two chromosomal abnormalities in the multi-probe biomarker sets were not demonstrated in the PBMCs.

Detection of CACs in control subjects may be due to the fact that some were heavy smokers and exposed to the same tobacco carcinogens as the cancer patients. The presence of chromosome damage in healthy controls at significantly lower levels than in cancer patients has been reported (42,43). Moreover there is evidence that abnormalities in cytogenetic biomarkers are positively correlated with cancer risk reflecting the genotoxic effects of carcinogens as well as individual susceptibility to cancer development (44–46).

In comparison with other methods that used EpCAM-positive cells to detect CTCs, we observed several orders of magnitude higher numbers of CACs in our antigen independent genetic assays than most other studies. Depending on the biomarker abnormality assayed, up to 45 CACs per micro liter were detected compared to < 10 CACs per 1 milliliter in most studies using immunomagnetic beads (2,6,15,17,41). The percentages of CACs tended to increase with disease stage reflecting tumor burden. Certain genetic abnormalities were associated with relapse and overall survival duration; however, after adjustment for age, sex, and disease stage these were not statistically significant most likely reflecting a relatively small number of patients and the strong effect of disease stage on overall survival.

CACs with certain chromosomal abnormalities were present in the blood of patients with both early and advanced stage NSCLC and were correlated with relapse and poor survival, suggesting that genetic abnormalities that persist throughout all stages and at relapse may represent cancer stem cell markers.

Although the primary lung tumors manifested a wide range of clonal heterogeneity, including subpopulations with high chromosomal copy number the usual phenotype for CACs in blood was a single loss or gain of a gene or chromosome which correlated with the presence of similar clones in the tumor (Table 6S). Our study is the first to show that high levels of CACs are present in the blood stream of patients with NSCLC according to detection of genetic abnormalities using antigen independent FISH-based assays. Our ongoing research shows that while most CACs are negative for CD45, a hemolymphoid marker, a low percentage of CACs are cytokeratin positive, and may co-express CD45. Future goals include studies with larger

numbers of patients in both early and advanced disease to validate that CACS are related to disease stage, relapse and survival, as well as evaluating epithelial, mesenchymal, stem cell and hematolymphoid markers, combined with FISH, in order to discover the origin and phenotype of the CACs. If, as we suspect, that CACs actually represent CTCs, then assays for CACs may be used for clinical applications. These would include use of CACs as an adjunct to the diagnosis of radiographically detected indeterminate lung nodules, or as markers of response to cancer therapy and for detection of minimal residual disease.

Translational Relevance

Circulating cytogenetically abnormal cells (CACs) containing similar genetic abnormalities to the primary tumor are present in patients with non-small cell lung cancer (NSCLC). We developed an antigen independent quantitative FISH-based assay that detected circulating CACs from patients with NSCLC using DNA probes to identify genomic aberrations known to be present in patients with NSCLC. Genetic abnormalities in CACs in baseline blood correlated with genetic abnormalities in resected NSCLC. Numbers of CACs correlated with stage and were significantly different from controls. Different from previous studies employing immuno-magnetic capture for epithelial antigens, with low tumor cell recovery, our approach allowed us to detect much higher numbers of circulating CACs, which may be epithelial, mesenchymal or stem cell in origin. Assays for CACs may be used as adjuncts to the diagnosis of indeterminate lung nodules, as markers of response to therapy and for detection of minimal residual disease.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

The authors wish to thank Mr. Jinping Zhang and Dr. Ying Chen for the preparation of in-house probes, Ms. Helen Yang for isolation and quantitation of mononuclear cells from blood specimens, Mr. Donald Norwood for editing the manuscript, Dr. J. Jack Lee for statistical review, and Ms. Cassandra Woodard for secretarial assistance.

This work was supported in part by the following grants: SPORE grant (#P50CA70907), a Cancer Prevention Fellowship funded by the National Cancer Institute, Grant K07CA093592, National Cancer Institute grants CA123208, CA129050 and CA55769, and in part by a grant from Astra Zeneca.

References

1. Fehm T, Sagalowsky A, Beitsch P, et al. Cytogenetic evidence that circulating epithelial cells in patients with carcinoma are malignant. *Clinical Cancer Research* July;2002 8:2073–2084. [PubMed: 12114406]
2. Maheswaran S, Sequist LV, Nagrath S, et al. Detection of mutations in EGFR in circulating lung-cancer cells. *N Engl J Med* 2008;359:366–377. [PubMed: 18596266]
3. Cristofanilli M, Budd GT, Ellis MJ, et al. Circulating tumor cells, disease progression, and survival in metastatic breast cancer. *N Engl J Med* 2004;351:781–791. [PubMed: 15317891]
4. Hayes DF, Cristofanilli M, Budd GT, et al. Circulating tumor cells at each follow-up time point during therapy of metastatic breast cancer patients predict progression-free and overall survival. *Clin Cancer Res* 2006;12:4218–4224. [PubMed: 16857794]
5. De Bono JS, Scher HI, Montgomery RB, et al. Circulating tumor cells predict survival benefit from treatment in metastatic castration-resistant prostate cancer. *Clin Cancer Res* 2008;14:6302–6309. [PubMed: 18829513]
6. Cohen SJ, Punt CJ, Iannotti N, et al. Relationship of circulating tumor cells to tumor response, progression-free survival, and overall survival in patients with metastatic colorectal cancer. *J Clin Oncol* 2008;26:3213–3221. [PubMed: 18591556]

7. Allard WJ, Matera J, Miller MC, et al. Tumor cells circulate in the peripheral blood of all major carcinomas but not in healthy subjects or patients with nonmalignant diseases. *Clin Cancer Res* 2004;10:6897–6904. [PubMed: 15501967]
8. Nagrath S, Sequist LV, Maheswaran S, et al. Isolation of rare circulating tumour cells in cancer patients by microchip technology. *Nature* 2007;450:1235–1239. [PubMed: 18097410]
9. Sieuwerts AM, Kraan J, Bolt J, et al. Anti-epithelial cell adhesion molecule antibodies and the detection of circulating normal-like breast tumor cells. *J Natl Cancer Inst* 2009;101:61–66. [PubMed: 19116383]
10. Zabaglo L, Ormerod MG, Parton M, et al. Cell filtration-laser scanning cytometry for the characterization of circulating breast cancer cells. *Cytometry A* 2003;55:102–108. [PubMed: 14505315]
11. Braun S, Naume B. Circulating and disseminated tumor cells. *J Clin Oncol* 2005;23:1623–1626. [PubMed: 15755968]
12. Wong SC, Chan CM, Ma BB, et al. Clinical significance of cytokeratin 20-positive circulating tumor cells detected by a refined immunomagnetic enrichment assay in colorectal cancer patients. *Clin Cancer Res* 2009;15:1005–1012. [PubMed: 19188172]
13. Pantel K, Brakenhoff RH, Brandt B. Detection, clinical relevance and specific biological properties of disseminating tumour cells. *Nat Rev Cancer* 2008;8:329–340. [PubMed: 18404148]
14. Paterlini-Brechot P, Benali NL. Circulating tumor cells (CTC) detection: Clinical impact and future directions. *Cancer Lett* 2007;253:180–204. [PubMed: 17314005]
15. Wu C, Hao H, Li L, et al. Preliminary investigation of the clinical significance of detecting circulating tumor cells enriched from lung cancer patients. *J Thorac Oncol* 2009;4:30–36. [PubMed: 19096303]
16. Gross HJ, Verwer B, Houck D, et al. Model study detecting breast cancer cells in peripheral blood mononuclear cells at frequencies as low as 10⁽⁻⁷⁾. *Proc Natl Acad Sci USA* 1995;92:537–541. [PubMed: 7831325]
17. Ntourogi TG, Ashraf SQ, McGregor SB, et al. Detection of circulating tumour cells in peripheral blood with an automated scanning fluorescence microscope. *Br J Cancer* 2008;99:789–795. [PubMed: 18682708]
18. Katz RL, Zaidi TM, Fernandez RL, et al. Automated detection of genetic abnormalities combined with cytology in sputum is a sensitive predictor of lung cancer. *Mod Pathol* 2008;21:950–960. [PubMed: 18500269]
19. Daniely M, Rona R, Kaplan T, et al. Combined analysis of morphology and fluorescence in situ hybridization significantly increases accuracy of bladder cancer detection in voided urine samples. *Urology* 2005;66:1354–1359. [PubMed: 16360483]
20. Jiang F, Yin Z, Caraway NP, et al. Genomic profiles in stage I primary non small cell lung cancer using comparative genomic hybridization analysis of cDNA microarrays. *Neoplasia* 2004;6:623–635. [PubMed: 15548372]
21. Li R, Todd NW, Qiu Q, et al. Genetic deletions in sputum as diagnostic markers for early detection of stage I non-small cell lung cancer. *Clin Cancer Res* 2007;13:482–487. [PubMed: 17255269]
22. Barkan GA, Caraway NP, Jiang F, et al. Comparison of molecular abnormalities in bronchial brushings and tumor touch preparations. *Cancer* 2005;105:35–43. [PubMed: 15605362]
23. Yendamuri S, Vaporciyan AA, Zaidi T, et al. 3p22.1 and 10q22.3 deletions detected by fluorescence in situ hybridization (FISH): a potential new tool for early detection of non-small cell lung Cancer (NSCLC). *J Thorac Oncol* 2008;3:979–984. [PubMed: 18758299]
24. Kholognyuk ID, Kost-Alimova M, Yang Y, et al. The microcell hybrid-based “elimination test” identifies a 1Mb putative tumor suppressor region at 3p22.2–p22.1 centromeric to the homozygous deletion region detected in lung cancer. *Genes Chromosomes Cancer* 2002 Jul;34(3):34–4.
25. Goeze A, Schluns K, Wolf G, Thasler Z, et al. Chromosomal imbalances of primary and metastatic lung adenocarcinomas. *J Pathol* 2002 Jan;196(1):8–16. [PubMed: 11748636]
26. Caballero OL, Cohen D, Liu Q, et al. Loss of chromosome arms 3p and 9p and inactivation of P16 (INK4a) in normal epithelium of patients with primary lung cancer. *Genes Chromosomes Cancer* 2001 Oct;32(2):119–25. [PubMed: 11550279]
27. Wistuba II, Behrens C, Virmani AK, et al. High resolution chromosome 3p allelotyping of human lung cancer and preneoplastic/preinvasive bronchial epithelium reveals multiple, discontinuous sites

- of 3p allele loss and three regions of frequent breakpoints. *Cancer Res* 2000 Apr 1;60(7):1949–60. [PubMed: 10766185]
28. Petersen S, Aninat-Meyer M, Schluns K, et al. Chromosomal alterations in the clonal evolution to the metastatic stage of squamous cell carcinomas of the lung. *Br J Cancer* 2000 Jan;82(1):65–73. [PubMed: 10638968]
 29. Shriver SP, Shriver MD, Tirpak DL, et al. Trinucleotide repeat length variation in the human ribosomal protein L14 gene (RPL14): localization to 3p21.3 and loss of heterozygosity in lung and oral cancers. *Mutat Res* 1998 Nov;406(1):9–23. [PubMed: 9920051]
 30. Massion PP, Zou Y, Uner H, et al. Recurrent genomic gains in preinvasive lesions as a biomarker of risk for lung cancer. *PLoS One* 2009;4(6):e5611. [PubMed: 19547694]
 31. Choi CM, Seo KW, Jang SJ, et al. Chromosomal instability is a risk factor for poor prognosis of adenocarcinoma of the lung: Fluorescence in situ hybridization analysis of paraffin-embedded tissue from Korean patients. *Lung Cancer* 2009 Apr;64(1):66–70. Epub 2008 Sep 23. [PubMed: 18814932]
 32. Kang JU, Koo SH, Kwon KC, Park JW, Jung SS. Gain of the EGFR gene located on 7p12 is a frequent and early event in squamous cell carcinoma of the lung. *Cancer Genet Cytogenet* 2008;184(1):31–7. [PubMed: 18558286]
 33. Kang JU, Koo SH, Kwon KC, park JW, Shin SY, Kim JM, Jung SS. High frequency of genetic alterations in non-small cell lung cancer detected by multi-target fluorescence in situ hybridization. *J Korean Med Sci* 2007 Sep;22(Suppl):S47–51. [PubMed: 17923754]
 34. Zudaire I, Lozano MD, Valquez MF, Pajares MJ, Agorreta J, Pio R, Zulueta JJ, Yankelevitz DF, Henschke CI, Montuenga LM. Molecular characterization of small peripheral lung tumor based on the analysis of fine needle aspirates. *Histol Histopathol* 2008 Jan;23(1):33–40. [PubMed: 17952855]
 35. Halling KC, Rickman OB, Kipp Br, Harwood AR, Doerr CH, Jett JR. A comparison of cytology and fluorescence in situ hybridization for the detection of lung cancer in bronchoscopic specimens. *Chest* 2006;130(3):694–701. [PubMed: 16963665]
 36. Flores-Staino C, Darai-Ramqvist E, Dobra K, Hjerpe A. Adaptation of a commercial fluorescent in situ hybridization test to the diagnosis of malignant cells in effusion. *Lung Cancer*. 2009 Jun 10;
 37. Aktas B, Tewes M, Fehm T, et al. Stem cell and epithelial-mesenchymal transition markers are frequently overexpressed in circulating tumor cells of metastatic breast cancer patients. *Breast Cancer Res* 2009;11(4):R46. Epub. 2009 Jul 9. [PubMed: 19589136]
 38. Jiang F, Qiu Q, Khanna A, Todd NW, Deepak J, Xing L, Wang H, Liu Z, Su Y, Stass SA, Katz RL. *Mol Cancer Res* 2009 Mar;7(3):330–8. Epub 2009 Mar 10. Aldehyde dehydrogenase 1 is a tumor stem cell-associated marker in lung cancer. [PubMed: 19276181]
 39. Dave BJ, Hopwood VL, King TM, Jiang H, Spitz MR, Pathak S. Genetic susceptibility to lung cancer as determined by lymphocytic chromosome analysis, Department of Cell Biology, University of Texas M. D. Anderson Cancer Center, Houston 77030, USA. *Cancer Epidemiol Biomarkers Prev* 1995 Oct–Nov;4(7):743–9. [PubMed: 8672991]
 40. Swennenhuis JF, Tibbe AG, Levink R, et al. Characterization of circulating tumor cells by fluorescence in situ hybridization. *Cytometry A* 2009;75(6):520–7. [PubMed: 19291800]
 41. Leversha MA, Han J, Asgari Z, et al. Fluorescence in situ hybridization analysis of circulating tumor cells in metastatic prostate cancer. *Clin Cancer Res* 2009;15(6):2091–7. Epub 2009. [PubMed: 19276271]
 42. El-Zein RA, Schabath MB, Etzel CJ, Lopez MS, Franklin JD, Spitz MR. Cytokinesis-blocked micronucleus assay as a novel biomarker for lung cancer risk. *Cancer Res* 2006;66(12):6449–56. [PubMed: 16778224]
 43. Varella-Garcia M, Chen L, Powell RL, Hirsch FR, Kennedy TC, Keith R, Miller YE, Mitchell JD, Franklin WA. Spectral karyotyping detects chromosome damage in bronchial cells of smokers and patients with cancer. *Am J Respir Crit Care Med* 2007;176(5):505–12. [PubMed: 17600274]
 44. Bonassi S, Znaor A, Norppa H, Hagmar L. Chromosomal aberrations and risk of cancer in humans: an epidemiologic perspective. *Cytogenet Genome Res* 2004;104:376–82. [PubMed: 15162068]
 45. Smerhovsky Z, Landa K, Rossner P, et al. Risk of cancer in an occupationally exposed cohort with increased level of chromosomal aberrations. *Environ Health Perspect* 2001;109:41–5. [PubMed: 11171523]

46. Tucker JD, Preston RJ. Chromosome aberrations, micronuclei, aneuploidy, sister chromatid exchanges, and cancer risk assessment. *Mutat Res* 1996;365:147–59. [PubMed: 8898995]





Figure 1.

Figure 1A: Calibration curves of actual versus expected recovery at different dilutions of spiked tumor cells showing polysomy (arrows) in a background of normal PBMCs (A) 3p22.1/CEP3 (B) 10q22.3/CEP10 (C) LAV (D) URO probes. X-axis depicts percentage of tumor cells recovered, Y-axis depicts cell dilutions from 0.05–0.0001 (X and Y-axis logarithmic scale).






Figure 2.

Figure 2A: Error Bar Plots Showing Percentage Deletions and Gains of EGFR (Y-axis) with the LAV probe set in PBMCs Specimens Obtained from Controls and Cases with NSCLC by Disease Stage (X-axis).

Figure 2B: Mean \pm SE CACs per μ l in controls and cases with NSCLC with chromosomal abnormalities of 3p22.1/CEP3, 10q22.3/CEP10, URO and LAV probes stratified by stage. Note the trend for numbers of CACs for all chromosomal abnormalities to increase from early to advanced stage of NSCLC.




Figure 3.

Figure 3A: Stage IA adenocarcinoma, (A, B) deletions 3p22.1; (C) deletions 10q22.3; (D) polysomy 10q22.3/CEP10.

Figure 3B: Stage IA adenocarcinoma, (A) monosomy 6p11-q12; (B) amplification EGFR, C-myc; (C) trisomy CEP3 and monosomy CEP17; and (D) polysomy CEP3, CEP7, CEP17, and 9p21.3.

Table 1

Demographic Characteristics of the Study Population

Characteristic	Controls N=24	Cases N=59	Clinical Stage				
			IA N=16	IB N=8	II N=7	III N=10	IV N=18
Gender: N (%) Male	9 (37.5)	30 (50.8)	11 (78.6)	4 (50.0)	2 (22.2)	5 (50.0)	8 (44.4)
P*		0.269					
Age (Mean±SE)	55.5±2.86	66.8±1.36	67.4±2.55	70.63±3.67	61.0±3.01	67.7±2.94	66.9±2.88
P*		0.001					
Ever Smokers (%)	45.83	89.83	81.25	75	100	90	100
P*		<0.001					
Pack Year (Mean±SE)	53.1±12.48	45.0±3.37	44.6±8.41	48.0±6.99	45.4±10.1	45.1±5.97	43.8±6.43
P*		0.702					
Histology: N (%)							
Squamous	12 (20.3)	6 (37.5)	2 (25.0)	2 (28.6)	1 (10.0)	1 (5.6)	
Adenocarcinoma	32 (54.2)	9 (64.3)	5 (62.5)	5 (71.4)	4 (40.0)	9 (50.0)	
NSC	15 (25.4)	1 (7.1)	1 (12.5)	0 (0.0)	5 (50.0)	8 (44.4)	
Patients who died: N (%)	22 (37.3)	1 (4.5)	0 (0.0)	0 (0.0)	6 (27.3)	15 (68.2)	

* P-value derived from Mann-Whitney test (continuous variables); Chi-square test for association (categorical variables); all P-values are two-sided and compare controls to all cases; NSC=Non-small Cell Carcinoma

Table 2A

Distribution of Biomarkers in Cytogenetically Abnormal Cells among Case and Controls

Marker % (Mean±SE)	Controls N=24	Cases N=59	P
deletion 3p22.1/CEP3	2.25±0.40	5.33±0.46	<0.001
gains 3p22.1/CEP3	0.17±0.07	0.79±0.11	<0.001
mono 3p22.1/CEP3	0.36±0.13	0.66±0.09	0.023
poly 3p22.1/CEP3	0.28±0.12	1.18±0.17	<0.001
Combined abnormalities	3.05±0.46	7.96±4.37	<0.001
deletion 10q22.3/CEP10	0.76±0.21	3.52±0.42	<0.001
gains 10q22.3/CEP10	0.43±0.11	1.20±0.18	0.003
mono 10q22.3/CEP10	0.47±0.16	1.11±0.14	0.005
poly 10q22.3/CEP 10	0.01±0.01	0.44±0.07	<0.001
Combined abnormalities	1.67±0.24	6.27±0.53	<0.001
3p22.1 and 10q22.3 Deletions	3.01±0.52	8.83±0.72	<0.001
LA Vysion Deletions			
Single	4.41±0.63	7.31±0.51	0.001
EGFR	0.77±0.12	1.54±0.18	0.007
5p15.2	0.45±0.17	1.85±0.27	<0.001
C-myc	0.15±0.06	0.64±0.10	0.002
6p11-q11	2.99±0.66	3.41±0.35	0.256
LA Vysion Gains			
Single	2.66±0.37	5.85±0.50	<0.001
EGFR	1.60±0.26	3.86±0.39	<0.001
5p15.2	0.35±0.12	0.68±0.12	0.1
C-myc	0.14±0.08	0.22±0.06	0.247
6p11-q11	0.65±0.16	1.02±0.14	0.18
All LA Vysion Abnormalities	0.65±0.22	2.18±0.26	<0.001
UroVysion Deletions			
Single	5.36±0.75	8.31±0.43	<0.001
CEP3	0.20±0.06	0.21±0.06	0.547
CEP7	0.15±0.06	1.06±0.18	<0.001
9p21.3	0.89±0.21	1.49±0.23	0.093
CEP 17	4.17±0.33	5.54±0.33	0.008
UroVysion Gains			
Single	2.23±0.31	5.35±0.42	<0.001
CEP3	1.22±0.19	3.26±0.32	<0.001
CEP7	0.52±0.14	0.73±0.13	0.466
9p21.3	0.23±0.10	0.64±0.10	0.003

Marker % (Mean±SE)	Controls N=24	Cases N=59	P
CEP17	0.26±0.07	0.64±0.10	0.047
All UroVysion Abnormalities	0.77±0.13	2.11±0.19	<0.001
URO and LAV % Abnormal	1.42±0.291	4.29±0.37	<0.001

P-value derived from Mann-Whitney test; all P-values are two-sided per *microliter.

Note: Enumeration of FISH Signals:

3p22.1 (green)/CEP 3 (red) or 10q22.3 (green)/CEP10 (red) - Normal: 2 signals for each probe; Deletion: Loss of one or both signals of 3p22.1 or 10q22.3 (green); Monosomy: Loss of signal for both probes (1red and 1green signal); Polysomy: more than 2 signals of each probe (3 red and 3 green or more); Gain: Gain of one or both signals of 3p22.1or 10q22.3 (green)

Table 2B
Cytogenetically Abnormal Cells by Biomarker Stratified by Pathological Stage of NSCLC expressed per Microliter of Peripheral Blood: Cases versus Controls

	CACs (Mean±SE)								
	Controls (Mean ± SE) N=24	IA (Mean ± SE) N=16	IB (Mean ± SE) N=8	II (Mean ± SE) N=7	III (Mean ± SE) N=10	IV (Mean ± SE) N=18	Cases (Mean ± SE) N=59	MW ^a	trend ^b
CAC per µl Peripheral Blood									
UroVysion	6.31±1.19	13.04±3.30	18.37±5.39	13.10±5.86	20.44±7.00	25.37±8.58	18.79±3.16	0.006	0.144
LAVysion	4.99±1.37	16.19±3.43	15.10±3.76	12.88±7.50	13.49±3.03	23.89±7.82	17.57±2.82	<0.001	0.349
3p22.1 Del/cep 3	7.04±2.81	43.75±10.58	17.28±6.87	27.09±11.95	34.94±10.35	71.12±19.82	45.52±7.49	0.011	0.13
Gain 3p22.1/cep 3	1.12±0.50	10.65±3.11	6.08±2.84	9.30±5.09	5.56±1.61	4.76±2.13	7.23±1.32	<0.001	0.094
10q22.3/cep 10 Del	5.58±1.44	20.72±16.25	25.74±7.90	18.15±6.87	20.86±4.93	35.98±8.52	25.78±3.28	<0.001	0.124
Gain 10q22.3/cep 10	3.44±0.95	8.46±1.70	7.08±3.19	6.52±2.33	14.22±7.92	12.38±5.06	10.21±2.12	0.013	0.328

^a P-value derived from Mann-Whitney test (all cases vs. controls);
^b P-value for trend of abnormality across tumor stage;
All P-values are two-sided (To express number per milliliter, multiply by 1000).
Abbreviations: NSCLC - Non-Small Cell Lung Cancer; CAC - Cytogenetically Abnormal Cells; URO - UroVysion; LAV - LAVysion; del - Deletion
Note: Enumeration of FISH Signals:

3p22.1 (green)/CEP 3 (red) or 10q22.3 (green)/CEP10 (red) - Normal: 2 signals for each probe; Deletion: Loss of one or both signals of 3p22.1 or 10q22.3 (green); Monosomy: Loss of signal for both probes (1 red and 1 green signal); Polysomy: more than 2 signals of each probe (3 red and 3 green or more); Gain: Gain of one or both signals of 3p22.1 or 10q22.3 (green)
LAV or URO Probes: Normal: 2 signals for each probe, Loss: Monosomy (one signal) of one probe; Gain: Polysomy (more than 2 signals) of one probe; Abnormal: Abnormality of two probes in one cell, loss or gain

Table 3

Markers associated with disease recurrence and Survival Recurrence

Marker	Median Survival (in months) 95% CI		P ^a	Cox Model Unadj HR (95% CI)	Cox Model Adjusted HR ^b (95% CI)
	Low	High			
3p22.1					
Deletions	--	18.8 (14.2–23.5)	0.071	--	--
Monosomy	90.5 (--)	18.9 (10.8–26.9)	0.024	--	--
Combined Abnormalities	--	18.9 (9.5–28.2)	0.03	--	--
3p/10q					
Deletions	29.0 (--)	15.5 (9.5–21.5)	0.019	--	--
10q22.3					
Monosomy	90.5 (--)	16.0 (9.7–22.3)	0.066	--	--
Combined Abnormalities	29.0 (10.7–47.3)	18.9 (9.9–27.8)	0.017	--	--
LAVysion					
EGFR del	--	18.9 (10.2–27.6)	0.034	--	--
6p del	15.5 (12.3–18.7)	90.4 (–)	0.01	--	--
Single gain	90.4 (–)	18.7 (14.0–23.7)	0.072	--	--
5p gain	29.0 (10.0–48.0)	16.0 (9.5–22.5)	0.046	3.04 (1.15–8.01)	1.57 (0.55–4.51)
6p gain	19.2 (7.6–30.8)	16.0 (8.9–23.1)	0.039	--	--
Abnormal	29.0 (14.8–43.2)	18.9 (10.9–26.8)	0.085	--	--
UroVysion					
CEP3 Deletions	38.4 (–)	16.0 (0.0–34.1)	0.072	4.03 (1.39–11.64)	1.87 (0.64–5.51)
9p21.3 Deletions	37.4 (0.0–80.8)	15.5 (12.4–18.6)	0.001	--	--
Single Gain	38.4 (4.2–72.6)	16.0 (10.1–22.0)	0.003	3.24 (1.13–9.26)	2.14 (0.67–6.79)
CEP3 Gain	38.4 (19.9–56.8)	16.0 (11.7–20.4)	0.004	--	--
Abnormal	29.0 (11.5–46.5)	18.9 (9.8–28.0)	0.028	--	--
Cytogenetically abnormal cells					
UroVysion	38.4 (10.6–66.1)	18.9 (7.8–30.0)	0.062	--	--

Marker	Median Survival (in months) 95% CI		P ^a	Cox Model Unadj HR (95% CI)	Cox Model Adjusted HR ^b (95% CI)
	Low	High			
URO/LAV	29.0 (11.0–47.0)	18.9 (10.3–27.4)	0.051	--	--
3p22.1	--	19.2 (7.1–31.3)	0.095	--	--
Survival					
10q22.3 Combined Abnormalities	29.0 (9.9–48.1)	19.2 (10.3–28.0)	0.071	--	--
3p22.1/10q22.3 Deletions	29.0 (--)	18.9 (14.2–23.5)	0.061	2.55 (1.02–6.38)	1.32 (0.47–3.72)
LAV EGFR Deletions	--	19.2 (9.5–28.9)	0.053	--	--
URO 9p21.3					
Deletions	28.4 (6.0–70.7)	19.2 (14.9–23.4)	0.054	--	--
Single Gain	38.4 (6.2–70.6)	18.9 (14.4–23.4)	0.015	3.62 (1.38–9.50)	2.51 (0.89–7.06)
CEP 3 Gain	29.0 (10.0–48.0)	18.9 (15.2–22.6)	0.027	--	--

P-value from Kaplan-Meier log-rank test, b HR adjusted by age, sex and stage; -- Median survival estimates not calculable.

Note: Enumeration of FISH Signals:

3p22.1 (green)/CEP 3 (red) or 10q22.3 (green)/CEP10 (red) - Normal: 2 signals for each probe; Deletion: Loss of one or both signals of 3p22.1 or 10q22.3 (green); Monosomy: Loss of signal for both probes (1 red and 1 green signal); Polysomy: more than 2 signals of each probe (3 red and 3 green or more); Gain: Gain of one or both signals of 3p22.1 or 10q22.3 (green)

LAV or URO Probes: Normal: 2 signals for each probe, Loss: Monosomy (one signal) of one probe; Gain: Polysomy (more than 2 signals) of one probe;